CANADIAN JOURNAL
OF BOTANY

JOURNAL CANADIEN
DE BOTANIQUE

VOLUME 65, 1987

Author Index / Index des auteurs

Aarssen, L.W., and Turkington, R. Responses to defoliation in *Holcus lanatus*, *Lolium perenne*, and *Trifolium repens* from three different-aged pastures, 1364
Abraham, D. see Etcheberry, R., 879
Acker, G. see Suske, J., 2098
Ackerley, C.A. see Rauser, W.E., 643
Ackerley, C.A. see Massicotte, H.B., 1127
Ackerley, C.A. see Massicotte, H.B., 1940
Agee, J.K., and Kertis, J. Forest types of the North Cascades National Park Service Complex, 1520
Ahmad, I., and Hellebust, J.A. Control of nitrogen assimilation in *Stichococcus bacillaris* by growth conditions, 432
Aist, J.R. see Smart, M.G., 745
Albertini, L. see Gélie, B., 2061
Aleksiuk, M.A. see Knowles, P., 1087
Alfieri, S.A., Jr. see El-Gholl, N.E., 1733
Al-Kherb, S.M., Roelfs, A.P., and Groth, J.V. Diversity for virulence in a sexually reproducing population of *Puccinia coronata*, 994
Allan, R. see Wallen, D.G., 1133
Al Maarri, K., Arnaud, Y., et Miginiac, E. Microbouturage *in vitro* de jeunes poiriers issus de pépins de ‹Passe Crassane›, 803
Ammar, S., Benbadis, A., and Tripathi, B.K. Floral induction in date palm seedlings (*Phoenix dactylifera* var. *Deglet Nour*) cultured *in vitro*, 137
Ammirati, J., Jr. see Carpenter, S.E., 716
Amundson, R.G. see Reich, P.B., 977
Anderson, A.J. see Griffiths, H.M., 63
Anderson, P.M. see Brubaker, L.B., 899
Andersson, D. see Solbreck, C., 2177
Andrus, R.E. see Wilcox, D.A., 2270
Anoruo, A.O. see Berlyn, G.P., 955
Antia, N.J. see Klut, M.E., 736
Ardouin, P., Jay, M., et Lumaret, R. Étude d'une situation de sympatrie entre diploïdes et tetraploïdes de *Dactylis glomerata* (Gramineae), sur la base du polymorphisme enzymatique et phénolique, 526
Armstrong, S.W., and Francis, D. Cell and nuclear area, RNA and protein content, and chromosome counts in meristematic cells of secondary roots and callus of *Cocos nucifera*, 547
Arnaud, Y. see Al Maarri, K., 803
Arora, D.K. see Filonow, A.B., 2084
Ashford, A.E. see Massicotte, H.B., 1927
Asselin, A. see Hogue, R., 476
Aston, M.J. see Hawkins, C.D.B., 666
Attiwill, P.M. see Maheswaran, J., 2601
Aujas, C. see Darmency, H., 2352
Babcock, C.E. see Shoemaker, R.A., 373
Babcock, C.E. see Shoemaker, R.A., 1921
Bailey, L.G. see Baum, B.R., 1563
Baker, D. see Newcomb, W., 80
Baker, D.M., Minor, H.C., and Cumbie, B.G. Scanning electron microscopy examination of soybean testa development, 2420
Bandoni, R.J. see Whitney, H.S., 95
Barabé, D. see Brouillet, L., 34
Barabé, D., Forget, S., et Chrétien, L. Organogénèse de la fleur de *Symplocarpus foetidus* (Araceae), 446
Barabé, D., Bergeron, Y., et Vincent, G. La répartition des caractères dans la classification des Hamamelididae (Angiospermae), 1756
Bärlocher, F. Aquatic hyphomycete spora in 10 streams of New Brunswick and Nova Scotia, 76
Barnes, B.V. see Denton, S.R., 779
Barnett, N.M. see Tabor, C.A., 1212
Barrasa, J.M. see Valmaseda, M., 1802
Barr, D.J.S., and Désaulniers, N.L. Ultrastructure of the *Lagena radicicola* zoospore, including a comparison with the primary and secondary *Saprolegnia* zoospores, 2161
Barrett, S.C.H., and Helenurm, K. The reproductive biology of boreal forest herbs. I. Breeding systems and pollination, 2036
Barrett, S.C.H. see Helenurm, K., 2047
Barron, G.L., and Thorn, R.G. Destruction of nematodes by species of *Pleurotus*, 774
Basappa, G.P., Muniyamma, M., and Chinnappa, C.C. An investigation of chromosome numbers in the genus *Brachiaria* (Poaceae: Paniceae) in relation to morphology and taxonomy, 2297

Bashan, Y., Okon, Y., and Henis, Y. Peroxidase, polyphenoloxidase, and phenols in relation to resistance against *Pseudomonas syringae* pv. *tomato* in tomato plants, 366
Basinger, J.F. see McIver, E.E., 2338
Baum, B.R. Classification of cultivated barley (*Hordeum vulgare*). 2. Elaboration of cultivar groups, 2152
Baum, B.R., and Bailey, L.G. A survey of endosperm starch granules in the genus *Hordeum*: a study using image analytic and numerical taxonomic techniques, 1563
Bayer, R.J. Morphometric analysis of western North American *Antennaria* (Asteraceae: Inuleae). I. Sexual species of sections *Alpinae*, *Dioicae*, and *Plantaginifoliae*, 2389
Bayer, R.J., La Duke, J.C., and Crawford, D. J. Isozyme variation in *Trillium nivale* (Liliaceae), 2250
Beamish, F.W.H., Jeffrey, K.A., Kolton, R.J., MacMahon, P.D., and Kowalchuk, K.A. Effects of the lampricide 3-trifluoromethyl-4-nitrophenol on diatoms in two southern Ontario streams, 893
Beck, C.B., and Stein, W.E., Jr. ***Galtiera bostonensis***, gen. et sp. nov., a protostelic calamopityacean from the New Albany shale of Kentucky, 348
Beck, R.C. see Berlyn, G.P., 955
Beckett, A., and Woods, A.M. The pattern of colony development and the formation of the uredinium of *Uromyces viciae-fabae* on *Vicia faba*, 1998
Beckett, A. see Woods, A.M., 2007
Bell, T.J., and Quinn, J.A. Effects of soil moisture and light intensity on the chasmogamous and cleistogamous components of reproductive effort of *Dichanthelium clandestinum* populations, 2243
Bellefleur, P. see Messier, C., 1682
Benbadis, A. see Ammar, S., 137
Bender, L. Fixation and metabolism of carbon dioxide by photosynthetically active carrot callus cultures, 1768
Benhamou, N., and Ouellette, G.B. Ultrastructural characterization of an extracellular fibrillar sheath on cells of *Ascocalyx abietina*, the scleroderris canker agent of conifers, 154
Benhamou, N., and Ouellette, G.B. Ultrastructural study and cytochemical investigation, by means of enzyme–gold complexes, of the fungus *Ascocalyx abietina*, 168
Benharrat, H., Renaudin, S., Rey, L., et Thalouarn, P. Sur la différenciation des trachéides dans les haustoriums d'*Osyris alba* parasitant *Hedera helix*, 1746
Bennett, K.D. Holocene history of forest trees in southern Ontario, 1792
Bérard-Therriault, L. see Poulin, M., 2686
Berendse, F., Berg, B., and Bosatta, E. The effect of lignin and nitrogen on the decomposition of litter in nutrient-poor ecosystems: a theoretical approach, 1116
Berg, B. see Berendse, F., 1116
Bergeron, Y. see Barabé, D., 1756
Berlyn, G.P., Anoruo, A.O., Beck, R.C., and Cheng, J. DNA content polymorphism and tissue culture regeneration in Caribbean pine, 955
Bernillon, D. see Gourbière, F., 131
Bertrand, C. see Brouillet, L., 34
Bhagat, N.R., Lalwani, H.B., and Reddy, P.S. Variation in node sequence in var. *hypogaea* collection of cultivated peanuts, 1245
Bidwell, R.G.S. see Trites, L.F., 1121
Biggs, A.R. see Rittinger, P.A., 1886
Bird, C.J. see Novaczek, I., 57
Birgersson, G. see Leufvén, A., 1038
Bisalputra, T. see Klut, M.E., 736
Bisson, M.A. see Hoffmann, R., 222
Black, L.D. see Warwick, S.I., 1396
Bliss, L.C. see Dawson, T.E., 863
Bliss, L.C. see Sohlberg, E.H., 1224
Blom, C.W.P.M. see Voesenek, L.A.C.J., 1638
Blouw, M.J. see Remphrey, W.R., 2188
Bocchieri, E., Floris, G., et Mulas, B. Dormance, germination et contenu protéique des semences de deux populations de *Schoenus nigricans* cueillies à des altitudes différentes, 617
Bock, C.A. see MacDonald, J.C., 703
Boelen, M.G. see Pryor, A., 2380
Boerner, R.E.J. see Dahlem, T.S., 242
Boerner, R.E.J. see Tyrrell, L.E., 1570
Bohm, L.R. see Maze, J., 1531
Boissière, J.-C., Boissière, M.-C., Champion-Arnaud, P., Lallemant, R., et Wagner, J. Le cycle des *Nostoc* des genres *Peltigera* et *Collema* en cultures *in vitro* et dans le thalle lichénique, 1468
Boissière, M.-C. see Boissière, J.-C., 1468
Bonfante-Fasolo, P., and Schubert, A. Spore wall architecture of *Glomus* spp., 539
Booth, C. see Reid, J., 1320
Booth, T. see Muhsin, T.M., 1137
Borden, J.H. see Brooks, J.E., 1249
Bosatta, E. see Berendse, F., 1116
Bosland, P.W., and Williams, P.H. An evaluation of *Fusarium oxysporum* from crucifers based on pathogenicity, isozyme polymorphism, vegetative compatibility, and geographic origin, 2067
Botha, F.C., and Dennis, D.T. Phosphoglyceromutase activity and concentration in the endosperm of developing and germinating *Ricinus communis* seeds, 1908
Bouchard, A. see Jean, M., 1969

Bourgeais, P., Guerrier, G., et Strullu, D.G. Adaptation au NaCl de *Lycopersicon esculentum* : étude comparative des cultures de cals ou de parties terminales de tiges, 1989
Bradfield, G.E. see Gagnon, D., 822
Brassard, G.R. see Etcheberry, R., 879
Bret, J.-P. see Robert, J.-C., 505
Brett, S.J. see Wetherbee, R., 1019
Briane, J.P. see Escarré, J., 2668
Bronner, R., Dreger-Jauffret, F., and Pfirsch, E. Lack of gravitropic response of tumour roots of *Forsythia intermedia*: cytological evidence for this behaviour, 143
Brooks, J.E., Borden, J.H., Pierce, H.D., Jr., and Lister, G.R. Seasonal variation in foliar and bud monoterpenes in Sitka spruce, 1249
Brouillet, L., Bertrand, C., Cuerrier, A., et Barabé, D. Les hydathodes des genres *Begonia* et *Hillebrandia* (Begoniaceae), 34
Brubaker, L.B., Graumlich, L.J., and Anderson, P.M. An evaluation of statistical techniques for discriminating *Picea glauca* from *Picea mariana* pollen in northern Alaska, 899
Buchheim, M.A., and Hoffman, L.R. Structure and reproduction of *Sphaeroplea fragilis* sp.nov., a new member of the Sphaeropleaceae (Chlorophyceae) from California, 2330
Budd, K., and Kerson, G.W. Uptake of phosphate by two cyanophytes: cation effects and energetics, 1901
Bunce, J.A. Species-specific responses to water stress of gas exchange parameters mimicked by applied abscisic acid, 103
Burgess, D., and Peterson, R.L. Development of *Alnus japonica* root nodules after inoculation with *Frankia* strain HFPArI$_3$, 1647
Burgess, D., and Peterson, R.L. Effect of nutrient conditions on root nodule development in *Alnus japonica*, 1658
Burzynski, M.P. see Wein, R.W., 1180
Cabot, E.L. see Druehl, L.D., 1599
Canne, J.M. Determinations of chromosome numbers in *Viola* (Violaceae), 653
Canne-Hilliker, J.M. see Kampny, C.M., 2255
Cardinal, A. see Poulin, M., 2686
Carlton, R.G., and Wetzel, R.G. Distributions and fates of oxygen in periphyton communities, 1031
Caron, G.E. see Owens, J.N., 1439
Carpenter, S.E., Trappe, J.M., and Ammirati, J., Jr. Observations of fungal succession in the Mount St. Helens devastation zone, 1980–1983, 716
Carpenter, S.E. see Seifert, K.A., 1262
Carr, G. see Mauperin, Ch., 2326
Catesson, A.-M. see Rao, K.S., 246
Catling, P.M., and Spicer, K.W. The perennial *Juncus* of section *Poiophylli* in the Canadian prairie provinces, 750
Cayouette, J. *Carex lyngbyei* excluded from the flora of eastern North America, and taxonomic notes on related species and hybrids, 1187
Chalifour, F.-P., and Nelson, L.M. Effects of continuous combined nitrogen supply on symbiotic dinitrogen fixation of faba bean and pea inoculated with different rhizobial isolates, 2542
Champion-Arnaud, P. see Boissière, J.-C., 1468
Chandra Sekhar, K.N., and Sawhney, V.K. Ontogenetic study of the fusion of floral organs in the normal and "solanifolia" mutant of tomato (*Lycopersicon esculentum*), 215
Chao, E.E., and Gruen, H.E. Intracellular activity of mycelial proteinases during fruit-body development in *Flammulina velutipes*, 518
Charlton, W.A. see Gould, K.S., 406
Chase, A.R. see El-Gholl, N.E., 1733
Checa, J., and Moreno, G. ***Leptosphaeria hispanica***, a new species of the Dothideales, Ascomycotina, 2096
Chen, T.H.H. see Gaudet, D.A., 1152
Cheng, J. see Berlyn, G.P., 955
Chinnappa, C.C. see Macdonald, S.E., 1726
Chinnappa, C.C. see Basappa, G.P., 2297
Chmielewski, J.G., and Ringius, G.S. Biomass-allocation patterns in populations of *Trillium erectum* and *T. grandiflorum* in southern Ontario, 1671
Chmielewski, J.G., Ringius, G.S., and Semple, J.C. The cytogeography of *Solidago uliginosa* (Compositae: Astereae) in the Great Lake region, 1045
Chmielewski, J.G. see Semple, J.C., 1047
Chmielewski, J.G. see Ringius, G.S., 2450
Chopin, T., and Floc'h, J.-Y. Seasonal variations of growth in the red alga *Chondrus crispus* on the Atlantic French coasts. I. A new approach by fluorescence labelling, 1014
Chrétien, L. see Barabé, D., 446
Christ, B.J., and Person, C.O. Effects of selection by host cultivars on populations of *Ustilago hordei*, 1379
Christ, B.J., and Person, C.O. Nonspecific action of polygenes in *Ustilago hordei*, 1390
Christ, B.J. see Person, C.O., 1384
Chuong, P.V. see Pauls, K.P., 834
Citharel, J. see Citharel, L., 1870
Citharel, L., et Citharel, J. Dualité des corps protéiques du mésophylle adaxial et abaxial des cotylédons de quelques espèces de la tribu des Génistées (Légumineuses), 1870
Clarkson, D.T. see Lefebvre, D.D., 1504
Clay, K. see Jones, J.P., 1027
Clifford, P.E., Offler, C.E., and Patrick, J.W. Injection of growth regulators into seeds growing *in situ* on plants of *Phaseolus vulgaris* with a double-fruit sink system, 612
Cline, M.L., France, R.C., and Reid, C.P.P. Intraspecific and interspecific growth variation of ectomycorrhizal fungi at different temperatures, 869
Cloutier, A. Microdistribution des espèces végétales sur des affleurements rocheux de dolomie, dans une érablière du sud du Québec, 1

Cole, K.M. see Tam, C.E., 532
Cole, K.M. see Oates, B.R., 1687
Coleman, J.S., Jones, C.G., and Smith, W.H. The effect of ozone on cottonwood – leaf rust interactions: independence of abiotic stress, genotype, and leaf ontogeny, 949
Cook, S.P., and Hain, F.P. Four parameters of the wound response of loblolly and shortleaf pines to inoculation with the blue-staining fungus associated with the southern pine beetle, 2403
Corlett, M. Erratum: *Pyxidiophora lundqvistii* n.sp. (Hypomycetales, Ascomycetes), 214
Cornu, A. see Vallade, J., 761
Correa, J. see Nielsen, R., 2467
Côté, R., Thain, J.F., and Fensom, D.S. Increase in electrical resistance of plasmodesmata of *Chara* induced by an applied pressure gradient across nodes, 509
Couchat, P. see Lascève, G., 1861
Couot-Gastelier, J., et Louguet, P. Étude comparée de la structure foliaire et de l'ultrastructure des stomates chez deux cultivars de *Pelargonium* × *hortorum*, l'un chlorophyllien et l'autre mutant à feuilles blanches, vertes ou panachées, 340
Coxson, D. Photoinhibition of net photosynthesis in *Stereocaulon virgatum* and *S. tomentosum*, a tropical – temperature comparison, 1707
Coxson, D.S., and Parkinson, D. The pattern of winter respiratory response to temperature, moisture, and freeze – thaw exposure in *Bouteloua gracilis* dominated grassland soils of southwestern Alberta, 1716
Crane, P.R., and Stockey, R.A. *Betula* leaves and reproductive structures from the Middle Eocene of British Columbia, Canada, 2490
Crawford, D.J. see Bayer, R. J., 2250
Cremers, G. see Sell, Y., 999
Cress, W.A., and Johnson, G.V. The effect of three osmotic agents on free proline and amino acid pools in *Atriplex canescens* and *Hilaria jamesii*, 799
Cuerrier, A. see Brouillet, L., 34
Cumbie, B.G. see Baker, D.M., 2420
Currah, R.S. ***Thanatephorus pennatus*** sp.nov. isolated from mycorrhizal roots of *Calypso bulbosa* (Orchidaceae) from Alberta, 1957
Currah, R.S., Sigler, L., and Hambleton, S. New records and new taxa of fungi from the mycorrhizae of terrestrial orchids of Alberta, 2473
Cusset, G. Erratum: La morphogenèse du limbe des Dicotylédones, 1550
Cutter, E.G. see Gould, K.S., 406
Czuba, M. see Weinberger, P., 696
Dahlem, T.S., and Boerner, R.E.J. Effects of canopy light gap and early emergence on the growth and reproduction of *Geranium maculatum*, 242
Darmency, H., and Aujas, C. Character inheritance and polymorphism in a wild oat (*Avena fatua*) population, 2352
David, A., and Rajchenberg, M. A reevaluation of *Wrightoporia* and *Amylonotus* (Aphyllophorales, Polyporaceae), 202
Davidson, C.G. see Remphrey, W.R., 2188
Dawson, T.E., and Bliss, L.C. Species patterns, edaphic characteristics, and plant water potential in a high-arctic brackish marsh, 863
DeChacin, C. see Weinberger, P., 696
Denford, K.E. see Downie, S.R., 559
Denford, K.E. see Fabijan, D.M., 2562
Dennis, D.T. see Botha, F.C., 1908
de Nooij, M.P., and van der Aa, H.A. *Phomopsis subordinaria* and associated stalk disease in natural populations of *Plantago lanceolata*, 2318
Denslow, J.S. Fruit removal rates from aggregated and isolated bushes of the red elderberry, *Sambucus pubens*, 1229
Denton, S.R., and Barnes, B.V. Application of the sucrose inversion method to delineation of region-wide temperature patterns, 779
Désaulniers, N.L. see Barr, D.J.S., 2161
Detlier, M.N. The distribution and reproductive phenology of intertidal fleshy crustose algae in Washington, 1838
Dorne, A.-J., Joyard, J., and Douce, R. Lipid metabolism of a Tertiary relic species, Kerguelen cabbage (*Pringlea antiscorbutica*), 2368
Douce, R. see Dorne, A.-J., 2368
Downie, S.R., and Denford, K.E. The biosystematics of *Arnica fulgens* and *A. sororia* (Asteraceae), 559
Dreger-Jauffret, F. see Bronner, R., 143
Druehl, L.D., Cabot, E.L., and Lloyd, K.E. Seasonal growth of *Laminaria groenlandica* as a function of plant age, 1599
Dubé, M., and Morisset, P. Morphological and leaf anatomical variation in *Festuca rubra sensu lato* (Poaceae) from eastern Quebec, 1065
Dubuc-Lebreux, M.-A., et Vieth J. Effets des rayons γ ^{60}Co sur des tigelles de *Gerbera jamesonii* irradiées in vitro, 261
Dubuc-Lebreux, M.-A., et Vieth, J. Effets de rayons γ ^{60}C sur des vitroplants haploïdes et diploïdes de *Gerbera jamesonii*, 2074
Ducreux, G. see Sihachakr, D., 192
Duke, S.H. see Hostak, M.S., 1108
Dumas, P., et Maillette, L. Rapport des sexes, effort et succès de reproduction chez *Rubus chamaemorus*, plant herbacée vivace dioïque de distribution subarctique, 2628
Duncan, D.R., and Himelick, E.B. The effects of amino acids on the growth and sporulation of *Verticillium dahliae*, 1299
Duncan, D.R. see Zehr, B.E., 491
Duysen, M.E. see Freeman, T.P., 2118
Ebba, T. see Person, C.O., 1384
Egger, K.N. see Turner, N.J., 921
Ehret, D.L. see Fisher, F.J.F., 2109
El-Gholl, N.E., Chase, A.R., Alfieri, S.A., Jr., and Schoulties, C.L. Occurrence of a microconidial state in *Cylindrocladium heptaseptatum*, 1733
El-Kassaby, Y.A. see Scagel, R.K., 1614
Elmhirst, J.F., and Heath, M.C. Interactions of the bean rust and cowpea rust fungi with species of the *Phaseolus–Vigna* plant complex. I. Fungal growth and development, 1096

Escarré, J., Houssard, C., et Briane, J.P. Évolution du rapport des sexes de populations de *Rumex acetosella* le long d'une succession postculturale, 2668
Etcheberry, R., Abraham, D., Brassard, G.R., et Favreau, M. Les mousses des îles St-Pierre-et-Miquelon, 879
Eversman, S., and Sigal, L.L. Effects of SO_2, O_3, and SO_2 and O_3 in combination on photosynthesis and ultrastructure of two lichen species, 1806
Faber-Langendoen, D., and Maycock, P.F. Composition and soil – environment analysis of prairies on Walpole Island, southwestern Ontario, 2410
Fabijan, D.M., Packer, J.G., and Denford, K.E. The taxonomy of the *Viola nuttallii* complex, 2562
Fabre, M.C. Étude de l'archégone de *Sphagnum palustre*. I. Le gamète femelle et la fécondation, 2233
Faria, S.M. de, McInroy, S.G., and Sprent, J.I. The occurrence of infected cells, with persistent infection threads, in legume root nodules, 553
Faris, M.A. see Ta, T.C., 2537
Favreau, M. see Etcheberry, R., 879
Felker, F.C., Peterson, D.M., and Nelson, O.E. Early grain development of the *seg2* maternal-effect shrunken-endosperm mutant of barley, 943
Fensom, D.S. see Côté, R., 509
Filonow, A.B., and Arora, D.K. Influence of soil matric potential on ^{14}C exudation from fungal propagules, 2084
Filtenborg, O. see Frisvad, J.C., 765
Fisher, F.J.F., Ehret, D.L., and Hollingdale, J. The pattern of vascular deployment near the pulvinus of the solar-tracking leaf of *Lavatera cretica* (Malvaceae), 2109
Fitter, A.H., and Parsons, W.F.J. Changes in phosphorus and nitrogen availability on recessional moraines of the Athabasca Glacier, Alberta, 210
Floc'h, J.-Y. see Chopin, T., 1014
Floris, G. see Bocchieri, E., 617
Foos, K.M., and Royer, J. Optimum growth temperature for *Pilobolus*, 1063
Forget, S. see Barabé, D., 446
Förster, H., and Mendgen, K. Immunocytochemical localization of pectinesterases in hyphae of *Phytophthora infestans*, 2607
Fortin, J.A. see Kropp, B.R., 500
Fowke, L.C. see Hakman, I., 656
France, R.C. see Cline, M.L., 869
Francis, D. see Armstrong, S.W., 547
Fraser, J.D. see Holthuijzen, A.M.A., 1092
Fredrick, L. see Staub, J.E., 792
Freeman, T.P., Duysen, M.E., and Williams, N.D. Effects of gene dosage of gene dosage on light harvesting chlorophyll accumulation, chloroplast development, and photosynthesis in wheat (*Triticum aestivum*), 2118
Frisvad, J.C., Filtenborg, O., and Wicklow, D.T. Terverticillate penicillia isolated from underground seed caches and cheek pouches of banner-tailed kangaroo rats (*Dipodomys spectabilis*), 765
Fromard, F. Systématique du taxon *Arctostaphylos uva-ursi*, Ericaceae, en Europe : données nouvelles concernant les populations pyrénéennes et circumpyrénéennes, 687
Fryer, G.I. see Johnson, E.A., 853
Funk, A. ***Monilia versiformia*** sp. nov. (Hyphomycetes) from a perennial Douglas-fir canker, 23
Funk, A. ***Stigmina zilleri*** sp. nov., associated with brown leaf spot of broadleaf maple, 482
Furnier, G.R. see Knowles, P., 1087
Gagnon, D., and Bradfield, G.E. Gradient analysis of west central Vancouver Island forests, 822
Gajewski, K. see Ritchie, J.C., 1605
Galen, C., and Plowright, R.C. Testing the accuracy of using peroxidase activity to indicate stigma receptivity, 107
Garbary, D.J. see Tam, C.E., 532
Garbaye, J. see Mauperin, Ch., 2326
Gaudet, D.A., and Kokko, E.G. Development of microsclerotia by *Coprinus psychromorbidus*, 26
Gaudet, D.A., and Chen, T.H.H. Effects of hardening and plant age on development of resistance to cottony snow mold (*Coprinus psychromorbidus*) in winter wheat under controlled conditions, 1152
Gaudet, D.A. see Traquair, J.A., 124
Gauthier, R. La répartition et l'habitat du *Tomenthypnum falcifolium* au Québec–Labrador, 286
Gélie, B., Petitprez, M., Souvre, A., et Albertini, L. Étude ultrastructurale des modifications induites par *Exserohilum turcicum* chez le limbe de *Zea mays* — Effet du gène de résistance *Ht-1*, 2061
Gignac, L.D. Ecological tolerance and niche structure of *Sphagnum* along a pollution gradient near Sudbury, Ontario, Canada, 1268
Given, D.R. see Whittier, D.P., 1770
Glime, J.M., and Vitt, D.H. A comparison of bryophyte species diversity and niche structure of montane streams and stream banks, 1824
Goates, B.J., and Hoffmann, J.A. Nuclear behavior during teliospore germination and sporidial development in *Tilletia caries*, *T. foetida*, and *T. controversa*, 512
Goudey, J.S., Saini, H.S., and Spencer, M.S. Seed germination of wild mustard (*Sinapis arvensis*): factors required to break primary dormancy, 849
Gould, K.S., Cutter, E.G., Young, J.P.W., and Charlton, W.A. Positional differences in size, morphology, and *in vitro* performance of pea axillary buds, 406
Gourbière, F., Pépin R., et Bernillon, D. Microscopie de la mycoflore des aiguilles de sapin (*Abies alba*). III. *Marasmius androsaceus*, 131
Grant, W.F. see MacLean, N.L., 1275
Graumlich, L.J. see Brubaker, L.B., 899
Gray, A.B., and Sackston, W.E. Host–parasite relationships in dual cultures of *Helianthus annuus* and *Plasmopara halstedii*, 2057
Gray, L.E. see Schechter, S.E., 1465

Green, D.G. Pollen evidence for the postglacial origins of Nova Scotia's forests, 1163
Greenwood, C.A. see Munn, E.A., 456
Grenville, D. see Malloch, D., 1281
Griffiths, H.M., and Anderson, A.J. Response of *Phaseolus vulgaris* protoplasts to chemical components of fungal origin, 63
Groth, J.V. see Al-Kherb, S.M., 994
Gruen, H.E. see Chao, E.E., 518
Guerrier, G. see Bourgeais, P., 1989
Gupta, B.K. see Kamal, 1259
Guries, R.P. see Wheeler, N.C., 1876
Hadden, K.A. see Ritchie, J.C., 1605
Hain, F.P. see Cook, S.P., 2403
Hakman, I., and Fowke, L.C. Somatic embryogenesis in *Picea glauca* (white spruce) and *Picea mariana* (black spruce), 656
Hambleton, S. see Currah, R.S., 2473
Hansen, E.M. see Hessburg, P.F., Sr., 962
Harmsen, R. see McBrien, H.L., 1478
Harris, J.M., and Singh, A.P. Unusual growth patterns in short shoots of *Pinus radiata*, 280
Harrison, L.G. see Lacalli, T.C., 1308
Hassall, M. see Visser, S., 635
Hattingh, M.J. see Mansvelt, E.L., 2517
Hattingh, M.J. see Mansvelt, E.L., 2523
Hawkins, C.D.B., Whitecross, M.I., and Aston, M.J. The effect of short-term aphid feeding on the partitioning of $^{14}CO_2$ photoassimilate in three legume species, 666
Hayes, D.C., and Seastedt, T.R. Root dynamics of tallgrass prairie in wet and dry years, 787
Heath, M.C. see Elmhirst, J.F., 1096
Helenurm, K., and Barrett, S.C.H. The reproductive biology of boreal forest herbs. II. Phenology of flowering and fruiting, 2047
Helenurm, K. see Barrett, S.C.H., 2036
Hellebust, J.A. see Ahmad, I., 432
Henis, Y. see Bashan, Y., 366
Henson, C.A. see Hostak, M.S., 1108
Hessburg, P.F., Sr., and Hansen, E.M. Pathological anatomy of black stain root disease of Douglas-fir, 962
Hicks, C.B., Stephens, L.C., and Weigle, J.L. *In vitro* pollen germination and viability of a Java × New Guinea *Impatiens* interspecific hybrid, 1967
Hicks, G.S. Adventitious rooting of apple microcuttings in vitro: an anatomical study, 1913
Hill, D.R.A. see Wetherbee, R., 1019
Himelick, E.B. see Duncan, D.R., 1299
Hinckley, T.M. see Schulte, P.J., 255
Hoffman, L.R. see Buchheim, M.A., 2330
Hoffmann, J.A. see Goates, B.J., 512
Hoffmann, R., and Bisson, M.A. *Chara buckellii*, a euryhaline charophyte from an unusual saline environment. II. Membrane potential and membrane conductance at steady state, 222
Hogue, R., and Asselin, A. Detection of 10 additional pathogenesis-related (*b*) proteins in intercellular fluid extracts from stressed 'Xanthi-nc' tobacco leaf tissue, 476
Hollingdale, J. see Fisher, F.J.F., 2109
Holthuijzen, A.M.A., Sharik, T.L., and Fraser, J.D. Dispersal of eastern red cedar (*Juniperus virginiana*) into pastures: an overview, 1092
Hopcroft, D.H. see Pankhurst, C.E., 2676
Hopkins, N.A. Mycorrhizae in a California serpentine grassland community, 484
Hostak, M.S., Henson, C.A., Duke, S.H., and VandenBosch, K.A. Starch-granule distribution between cell types of alfalfa nodules as affected by symbiotic development, 1108
Houssard, C. see Escarré, J., 2668
Howarth, M. see Kott, L.S., 53
Howes, N.K. see Kim, W.K., 1078
Howes, N.K. see Kim, W.K., 1785
Huang, H.C., and Kokko, E.G. Ultrastructure of hyperparasitism of *Coniothyrium minitans* on sclerotia of *Sclerotinia sclerotiorum*, 2483
Hubart, J.-M. see Malloch, D., 1281
Hubart, J.-M. see Malloch, D., 2384
Hume, L. Long-term effects of 2,4-D application on plants. I. Effects on the weed community in a wheat crop, 2530
Huner, N.P.A. see Newsted, W.J., 233
Hunt, G.A., and Trappe, J.M. Seasonal hypogeous sporocarp production in a western Oregon Douglas-fir stand, 438
Ireland, R.R., and Spence, J.R. ***Rhacomitrium pacificum***, a new moss species from western North America, 859
Israel, H.W. see Smart, M.G., 745
Jana, S., and Thai, K.M. Patterns of changes of dormant genotypes in *Avena fatua* populations under different agricultural conditions, 1741
Janex-Favre, M.C. see Parguey-Leduc, A., 1491
Järlfors, U. see Shain, L., 1586
Järlfors, U. see Siegel, M.R., 2357
Jay, M. see Ardouin, P., 526
Jean, M., et Bouchard, A. La végétation de deux tourbières de la municipalité régionale de comté du Haut-Saint-Laurent (Québec), 1969
Jefferies, R.L. see Wolff, S.L., 1410
Jefferies, R.L. see Wolff, S.L., 1420
Jeffrey, K.A. see Beamish, F.W.H., 893

Johnson, E.A., and Fryer, G.I. Historical vegetation change in the Kananaskis Valley, Canadian Rockies, 853
Johnson, F.D. see Kramer, N.B., 1961
Johnson, G.V. see Cress, W.A., 799
Johnson, M.C. see Siegel, M.R., 2357
Johnson, R.G., Jones, E.B.G., and Moss, S.T. Taxonomic studies of the Halosphaeriaceae: *Ceriosporopsis*, *Haligena*, and ***Appendichordella*** gen.nov., 931
Johnson, W.C., Sharik, T.L., Mayes, R.A., and Smith, E.P. Nature and cause of zonation discreteness around glacial prairie marshes, 1622
Jolliffe, P.A. see Kristie, D.N., 2017
Jones, C.G. see Coleman, J.S., 949
Jones, E., and Wheelwright, N.T. Seasonal changes in the fruits of *Viburnum opulus*, a fleshy-fruited temperate-zone shrub, 2291
Jones, E.B.G. see Johnson, R.G., 931
Jones, J.P., and Clay, K. Ascus and crozier development in the Balansiae, 1027
Jones, W.T. see Pankhurst, C.E., 2676
Joyard, J. see Dorne, A.-J., 2368
Kamal, Gupta, B.K., and Verma, R.K. New species of *Cercospora* and *Cercosporidium* from Uttar Pradesh, India, 1259
Kammereck, R. see Ketchie, D.O., 2640
Kampny, C.M., and Canne-Hilliker, J.M. Patterns of floral development in *Agalinis* and allies (Scrophulariaceae). I. Floral development of *Agalinis fasciculata* and *A. tenuifolia*, 2255
Kasha, K.J. see Kott, L.S., 53
Kato, M., and Shimizu, S. Chlorophyll metabolism in higher plants. VII. Chlorophyll degradation in senescing tobacco leaves; phenolic-dependent peroxidative degradation, 729
Kausch, A.P. see Smereka, K.J., 2549
Kawchuk, L.M., and Nielsen, J. Improved bunt inoculation of wheat utilizing adventitious shoots, 1284
Kenkel, N.C. Trends and interrelationships in boreal wetland vegetation, 12
Kerson, G.W. see Budd, K., 1901
Kertis, J. see Agee, J.K., 1520
Ketchie, D.O., and Kammereck, R. Seasonal variation of cold resistance in *Malus* woody tissue as determined by differential thermal analysis and viability tests, 2640
Kikuzawa, K. Development and survival of leaves in *Magnolia obovata* in a deciduous broad-leaved forest in Hokkaido, northern Japan, 412
Kim, W.K., Howes, N.K., and Martens, J.W. Evidence for relatedness between *Puccinia graminis* f.sp. *secalis* and *P. graminis* f.sp. *tritici* from two-dimensional polypeptide mapping, 1078
Kim, W.K., and Howes, N.K. Localization of glycopeptides and race-variable polypeptides in urediosporelings and urediosporeling walls of *Puccinia graminis tritici*; affinity to concanavalin A, soybean agglutinin, and *Lotus* lectin, 1785
King, J. see Saxena, P.K., 1736
Klut, M.E., Bisalputra, T., and Antia, N.J. Some observations on the structure and function of the dinoflagellate pusule, 736
Knowles, P., Furnier, G.R., Aleksiuk, M.A., and Perry, D.J. Significant levels of self-fertilization in natural populations of tamarack, 1087
Kohlmeyer, J., and Volkmann-Kohlmeyer, B. Marine fungi from Aldabra, the Galapagos, and other tropical islands, 571
Kohmoto, K. see Park, P., 330
Kokko, E.G. see Gaudet, D.A., 26
Kokko, E.G. see Traquair, J.A., 124
Kokko, E.G. see Huang, H.C., 2483
Kolton, R.J. see Beamish, F.W.H., 893
Kott, E. see Kott, L.S., 53
Kott, L.S., Kott, E., Howarth, M., and Kasha, K.J. A comparative study of initiation and development of embryogenic callus from haploid embryos of several barley cultivars. I. Developmental state of embryo explants and callus potential, 53
Kowalchuk, K.A. see Beamish, F.W.H., 893
Kramer, N.B., and Johnson, F.D. Mature forest seed banks of three habitat types in central Idaho, 1961
Kristie, D.N., and Jolliffe, P.A. A rapid phytochrome-mediated growth response in etiolated *Sinapis alba* hypocotyls, 2017
Kropp, B.R., McAfee, B.J., and Fortin, J.A. Variable loss of ectomycorrhizal ability in monokaryotic and dikaryotic cultures of *Laccaria bicolor*, 500
Kuhnlein, H.V. see Turner, N.J., 921
Lacalli, T.C., and Harrison, L.G. Turing's model and branching tip growth: relation of time and spatial scales in morphogenesis, with application to *Micrasterias*, 1308
Lachaud, S. Xylogénèse chez les Dicotylédones arborescentes. V. Formation du bois de tension et transport de l'acide indole acétique tritié chez le Hêtre, 1253
La Duke, J.C. see Bayer, R.J., 2250
Laliberté, S., et Vieth, J. Modifications structurales des faisceaux vasculaires primaires de la tige de *Cordyline indivisa* (Agavaceae). 1. Analyse du patron de conversion, 299
Laliberté, S., et Vieth, J. Modifications structurales des faisceaux vasculaires primaires de la tige de *Cordyline indivisa* (Agavaceae). 2. Répartition des différents stades structuraux dans le cylindre primaire, en fonction de la distance sous l'apex, 324
Lallemant, R. see Boissière, J.-C., 1468
Lalonde, M. see St-Laurent, L., 1356
Lalwani, H.B. see Bhagat, N.R., 1245
Landré, P. see Nougarède, A., 907
Larkin-Lieffers, P.A. see Lieffers, V.J., 1371
Larson, D.W. see Matthes-Sears, U., 182
Lascève, G., Vavasseur, A., Couchat, P., Pellegrini, L., et Pellegrini, M. Ultrastructure des mitochondries des stomates de maïs; cas d'une ouverture photoactive, 1861
Latch, G.C.M. see Siegel, M.R., 2357

Law, S.K. see Yeung, E.C., 1459
Lechowicz, M.J. see Melancon, S., 1157
Lee, P.F. Ecological relationships of wild rice, *Zizania aquatica*. 5. Enhancement of wild rice production by *Potamogeton robbinsii*, 1433
Lee, P.F. see Stevenson, S.C., 2128
Lefebvre, D.D., and Clarkson, D.T. High-affinity phosphate absorption is independent of the root cell wall in *Pisum sativum*, 1504
Leone, G., and Van den Heuvel, J. Regulation by carbohydrates of the sequential *in vitro* production of pectic enzymes by *Botrytis cinerea*, 2133
Le Tacon, F. see Mauperin, Ch., 2326
Leufvén, A., and Birgersson, G. Quantitative variation of different monoterpenes around galleries of *Ips typographus* (Colleoptera: Scolytidae) attacking Norway spruce, 1038
Leung, H., and Williams, P.H. Nuclear division and chromosome behavior during meiosis and ascosporogenesis in *Pyricularia oryzae*, 112
Lieffers, V.J., and Rothwell, R.L. Rooting of peatland black spruce and tamarack in relation to depth of water table, 817
Lieffers, V.J., and Larkin-Lieffers, P.A. Slope, aspect, and slope position as factors controlling grassland communities in the coulees of the Oldman River, Alberta, 1371
Lindstrom, S.C., and Scagel, R.F. The marine algae of British Columbia, northern Washington, and southeast Alaska: division Rhodophyta (red algae), class Rhodophyceae, order Gigartinales, family Dumontiaceae, with an introduction to the order Gigartinales, 2202
Lister, G.R. see Brooks, J.E., 1249
Lloyd, K.E. see Druehl, L.D., 1599
Lloyd, R.M. see Warne, T.R., 362
Lobban, C.S., and Mann, D.G. The systematics of the tube-dwelling diatrom *Nitzschia martiana* and *Nitzschia* section *Spathulatae*, 2396
Louguet, P. see Couot-Gastelier, J., 340
Loza-Tavera, H., Serrano, B., Molina, J.D., Ortega-Delgado, M.L., and Sanchez-de-Jiménez, E. CO_2-fixation enzymes in maize (*Zea mays*) and grain yield, 607
Lumaret, R. see Ardouin, P., 526
Lundborg, A. Two antagonistic fungi, "D 37" and *Scytalidium album*, reduce the formation of bore holes by *Heterobasidion annosum* in Norway spruce (*Picea abies*), 916
Luttrell, E.S. Relations of hyphae to host cells in smut galls caused by species of *Tilletia*, *Tolyposporium*, and *Ustilago*, 2581
Lyrene, P.M. see Vander Kloet, S.P., 660
Macdonald, A.D., and Mothersill, D.H. Shoot development in *Betula papyrifera*. VI. Development of the reproductive structures, 466
MacDonald, J.C., and Bock, C.A. Spectral properties of pseudobactin produced by *Pseudomonas* sp. B10, 703
Macdonald, P. see Sinclair, J., 2147
Macdonald, S.E., Chinnappa, C.C., Reid, D.M., and Purdy, B.G. Population differentiation of the *Stellaria longipes* complex within Saskatchewan's Athabasca sand dunes, 1726
Macdowall, F.D.H. see Ta, T.C., 2537
Machardy, W.E. see Smereka, K.J., 2549
MacLean, N.L., and Grant, W.F. Evaluation of birdsfoot-trefoil (*Lotus corniculatus*) regenerated plants following *in virto* selection for herbicide tolerance, 1275
MacMahon, P.D. see Beamish, F.W.H., 893
Maekawa, N. A new species of the genus *Cerinomyces*, 583
Maheswaran, J., and Attiwill, P.M. Loss of organic matter, elements, and organic fractions in decomposing *Eucalyptus microcarpa* leaf litter, 2601
Maillette, L. see Dumas, P., 2628
Maizonnier, D. see Vallade, J., 761
Majumdar, S.K. see Wasserman, J.L., 2622
Malek, L. Protein mobilization from cotyledons of de-shooted pea seedlings, 1083
Malloch, D., and Hubart, J.-M. An undescribed species of *Microascus* from the Cave of Ramioul, 2384
Malloch, D., Grenville, D., and Hubart, J.-M. An unusual subterranean occurrence of fossil fungal sclerotia, 1281
Mann, D.G. see Lobban, C.S., 2396
Mansvelt, E.L., and Hattingh, M.J. Scanning electron microscopy of colonization of pear leaves by *Pseudomonas syringae* pv. *syringae*, 2517
Mansvelt, E.L., and Hattingh, M.J. Scanning electron microscopy of pear blossom invasion by *Pseudomonas syringae* pv. *syringae*, 2523
Martens, J.W. see Kim, W.K., 1078
Martínez, A.T. see Valmaseda, M., 1802
Marty, T.L. see Staub, J.E., 792
Massicotte, H.B., Ackerley, C.A., and Peterson, R.L. Localization of three sugar residues in the interface of ectomycorrhizae synthesized between *Alnus crispa* and *Alpova diplophloeus* as demonstrated by lectin binding, 1127
Massicotte, H.B., Peterson, R.L., and Ashford, A.E. Ontogeny of *Eucalyptus pilularis – Pisolithus tinctorius* ectomycorrhizae. I. Light microscopy and scanning electron microscopy, 1927
Massicotte, H.B., Peterson, R.L., and Ackerley, C.A. Ontogeny of *Eucalyptus pilularis – Pisolithus tinctorius* ectomycorrhizae. II. Transmission electron microscopy, 1940
Mathewes, R.W. see Wainman, N., 2179
Matthes-Sears, U., Nash, T.H. III, and Larson, D.W. The ecology of *Ramalina menziesii*. VI. Laboratory responses of net CO_2 exchange to moisture, temperature, and light, 182
Mauperin, Ch., Mortier, F., Garbaye, J., Le Tacon, F., and Carr, G. Viability of an ectomycorrhizal inoculum produced in a liquid medium and entrapped in a calcium alginate gel, 2326
Maycock, P.F. see Faber-Langendoen, D., 2410
Mayes, R.A. see Johnson, W.C., 1622
Maze, J., Scagel, R., and Bohm, L.R. Quantitative studies in ovule development. III. An estimate of shape changes in *Phyllostachys aurea*, 1531
Maze, J. see Scagel, R.K., 1614

McAfee, B.J. see Kropp, B.R., 500
McBrien, H.L., and Harmsen, R. Growth response of goldenrod, *Solidago canadensis* (Asteraceae), to periodic defoliation, 1478
McGraw, J.B. Seed-bank properties of an Appalachian sphagnum bog and a model of the depth distribution of viable seeds, 2028
McInroy, S.G. see Faria, S.M. de, 553
McIntyre, G.I. Inhibition by carbon dioxide of the phototropic response of the *Avena* coleoptile to blue light, 488
McIntyre, G.I. The role of water in the regulation of plant development, 1287
McIntyre, G.I. Studies on the growth and development of *Agropyron repens*: interacting effects of humidity, calcium, and nitrogen on growth of the rhizome apex and lateral buds, 1427
McIver, E.E., and Basinger, J.F. ***Mesocyparis borealis*** gen. et. sp. nov.: fossil Cupressaceae from the early Tertiary of Saskatchewan, Canada, 2338
McLachlan, J. see Novaczek, I., 57
McNeill, J. see Wolf, S.J., 647
Melancon, S., and Lechowicz, M.J. Differences in the damage caused by glaze ice on codominant *Acer saccharum* and *Fagus grandifolia*, 1157
Meletti, P. see Stefani, A., 1948
Mendgen, K. see Förster, H., 2607
Mercier, J., and Reeleder, R.D. Interactions between *Sclerotinia sclerotiorum* and other fungi on the phylloplane of lettuce, 1633
Messier, C., et Bellefleur, P. Variations des propriétés spectrales des feuilles au cours de la croissance du bouleau jaune, 1682
Miginiac, E. see Al Maarri, K., 803
Miller, P.M. Physiological responses of *Vaccinium vitis-idaea* to high tissue concentrations of manganese, 1643
Milligan, M.J., and Williams, P.G. Orchidaceous rhizoctonias from roots of nonorchids: mycelial and cultural characteristics of field and pot culture isolates, 598
Mims, C.W., Richardson, E.A., and Roberson, R.W. Ultrastructure of basidium and basidiospore development in three species of the fungus *Exobasidium*, 1236
Mineo, L. see Wasserman, J.L., 2622
Minor, H.C. see Baker, D.M., 2420
Mitchell, J.K., Smith, D.H., and Taber, R.A. Potential for biological control of *Cercosporidium personatum* leafspot of peanuts by *Dicyma pulvinata*, 2263
Molina, J.D. see Loza-Tavera, H., 607
Montant, C. see Parguey-Leduc, A., 1491
Mooney, P.A., and Van Staden, J. Induction of embryogenesis in callus from immature embryos of *Persea americana*, 622
Moore-Landecker, E. Effects of medium composition and light on formation of apothecia and sclerotia by *Pyronema domesticum*, 2276
Moreno, G. see Checa, J., 2096
Morisset, P. see Dubé, M., 1065
Mortier, F. see Mauperin, Ch., 2326
Moss, S.T. see Johnson, R.G., 931
Mothersill, D.H. see Macdonald, A.D., 466
Mozafar, A., and Oertli, J.J. Effect of temperature on ion efflux properties of leaf cell compartments in several chilling-sensitive and chilling-insensitive crop plants, 1893
Muhsin, T.M., and Booth, T. Fungi associated with halophytes of an inland salt marsh, Manitoba, Canada, 1137
Mulas, B. see Bocchieri, E., 617
Muniyamma, M. see Basappa, G.P., 2297
Munn, E.A., Greenwood, C.A., and Orpin, C.G. Organization of the kinetosomes and associated structures of zoospores of the rumen chytridiomycete *Neocallimastix patriciarum*, 456
Murrin, F., and Nolan, R.A. Ultrastructure of the infection of spruce budworm larvae by the fungus *Entomophaga aulicae*, 1694
Muruganandam, V. see Suryanarayanan, T.S., 815
Nagasawa, E. see Redhead, S.A., 972
Nakamura, R.R., and Stanton, M.L. Cryptic seed abortion and the estimation of ovule fertilization, 2463
Nash, T.H. III see Matthes-Sears, U., 182
Neish, G.A. Neotypification of *Fusarium tricinctum*, 589
Nelson, L.M. see Chalifour, F.P., 2542
Nelson, O.E. see Felker, F.C., 943
Newcomb, W., Baker, D., and Torrey, J.G. Ontogeny and fine structure of effective root nodules of the autumn olive (*Elaeagnus umbellata*), 80
Newmeyer, D. see Raju, N.B., 1539
Newsted, W.J., and Huner, N.P.A. Major polypeptides associated with differentiation in psychrophilic fungi, 233
Nicholls, K.H. Form variation in *Mallomonas asmundiae* and a description of ***Mallomonas sphagniphila*** sp.nov. (series *Corncnticae*, Mallomonadaceae), 627
Nicholls, K.H. The distinction between *Mallomonas acaroides* var. *acaroides* and ***Mallomonas acaroides*** var. ***muskokana*** var.nov. (Chrysophyceae), 1779
Nielsen, J. Reaction of *Hordeum* species to the smut fungi *Ustilago nuda* and *U. tritici*, 2024
Nielsen, J. see Kawchuk, L.M., 1284
Nielsen, R., and Correa, J. A comparative study of *Gomontia polyrhiza* and ***Chlorojackia pachyclados*** gen. et sp. nov. (Chlorophyta), 2467
Nishimura, S. see Park, P., 330
Nolan, R.A. see Murrin, F., 1694
Nougarède, A., Rondet, P., Landré, P., et Rembur, J. Effet d'un traitement par l'acide abscisique sur la division cellulaire, les teneurs en ADN et l'élongation du bourgeon cotylédonaire de plants de pois décapités, 907
Novaczek, I., Bird, C.J., and McLachlan, J. Phenology and temperature tolerance of the red algae *Chondria baileyana*, *Lomentaria baileyana*, *Griffithsia globifera*, and *Dasya baillouviana* in Nova Scotia, 57

Novaczek, I. Periodicity of epiphytes on *Zostera marina* in two embayments of the southern Gulf of St. Lawrence, 1676
Oates, B.R., and Cole, K.M. Ultrastructure and ontogeny of hair filaments in the subtidal saccate alga *Hydroclathrus clathratus* (Phaeophyta, Scytosiphonales), 1687
Oberwinkler, F. see Whitney, H.S., 95
Oertli, J.J. see Mozafar, A., 1893
Offler, C.E. see Clifford, P.E., 612
Ogden, J.G., III Vegetational and climatic history of Nova Scotia. I. Radiocarbon-dated pollen profiles from Halifax, Nova Scotia, 1482
Ohno, T. see Park, P., 330
Okon, Y. see Bashan, Y., 366
Olesen, J.M. Heterostyly, homostyly, and long-distance dispersal of *Menyanthes trifoliata* to Greenland, 1509
Olson, A.R. Ovule maturity in relation to blossom age of a diploid clone of *Solanum tuberosum*, 706
Onnis, A. see Stefani, A., 1948
Orpin, C.G. see Munn, E.A., 456
Ortega-Delgado, M.L. see Loza-Tavera, H., 607
Otani, H. see Park, P., 330
Ouellette, G.B. see Benhamou, N., 154
Ouellette, G.B. see Benhamou, N., 168
Owens, J.N., Simpson, S.J., and Caron, G.E. The pollination mechanism of Engelmann spruce (*Picea engelmannii*), 1439
Packer, J.G. see Fabijan, D.M., 2562
Paine, T.D., and Stephen, F.M. The relationship of tree height and crown class to the induced plant defenses of loblolly pine, 2090
Paine, T.D., and Stephen, F.M. Response of loblolly pine to different inoculum doses of *Ceratocystis minor*, a blue-stain fungus associated with *Dentroctonus frontalis*, 2093
Pankhurst, C.E., Hopcroft, D.E., and Jones, W.T. Comparative morphology and flavolan content of *Rhizobium loti* induced effective and ineffective root nodules on *Lotus* species, *Leuceana leucocephala*, *Carmichaelia flagelliformis*, *Ornithopus sativus*, and *Clianthus puniceus*, 2676
Parguey-Leduc, A., Janex-Favre, M.C., et Montant, C. Formation et évolution des ascospores de *Tuber melanosporum* (truffe noire du Périgord, Discomycètes), 1491
Parkinson, D. see Visser, S., 635
Parkinson, D. see Coxson, D.S., 1716
Park, P., Ohno, T., Nishimura, S., Kohmoto, K., and Otani, H. Leakage of sodium ions from plasma membrane modification, associated with permeability change, in host cells treated with a host-specific toxin from a Japanese pear pathotype of *Alternaria alternata*, 330
Parsons, W.F.J. see Fitter, A.H., 210
Patrick, J.W. see Clifford, P.E., 612
Patton, R.F. see Stanosz, G.R., 2124
Pauls, K.P., and Chuong, P.V. Flow cytometric identification of *Brassica napus* protoplast fusion products, 834
Peabody, D.C. see Peabody, R.B., 69
Peabody, R.B., and Peabody, D.C. Haploid monokaryotic basidiocarp tissues in species of *Armillaria*, 69
Pearcy, R.W. see Vallejos, C.E., 1303
Peirson, D.R. see Rittinger, P.A., 1886
Pellegrini, L. see Lascève, G., 1861
Pellegrini, M. see Lascève, G., 1861
Pépin R. see Gourbière, F., 131
Perkins, D.D. see Raju, N.B., 1539
Peron, J.Y., and Regnier, E. In vitro propagation of *Crambe maritima*, 72
Perry, D.J. see Knowles, P., 1087
Person, C.O., Christ, B.J., and Ebba, T. Polygenic modification of a major virulence gene in *Ustilago hordei*, 1384
Person, C.O. see Christ, B.J., 1379
Person, C.O. see Christ, B.J., 1390
Peterson, D.M. see Felker, F.C., 943
Peterson, R.L. see Massicotte, H.B., 1127
Peterson, R.L. see Burgess, D., 1647
Peterson, R.L. see Burgess, D., 1658
Peterson, R.L. see Massicotte, H.B., 1927
Peterson, R.L. see Massicotte, H.B., 1940
Petitprez, M. see Gélie, B., 2061
Pfirsch, E. see Bronner, R., 143
Phipps, J.B. see Smith, P.G., 2646
Pierce, H.D., Jr. see Brooks, J.E., 1249
Pizzolato, T.D. Vascular system of the floret of *Alopecurus carolinicnus* (Gramineae), 2592
Ploetz, R.C., and Shokes, F.M. Infection of different plant parts of soybean seedlings by southern *Diaporthe phaseolorum* and its role in the development of stem canker symptoms, 2104
Plowright, R.C. Corolla depth and nectar concentration: an experimental study, 1011
Plowright, R.C. see Galen, C., 107
Polowick, P.L., and Sawhney, V.K. A scanning electron microscopic study on the influence of temperature on the expression of cytoplasmic male sterility in *Brassica napus*, 807
Poulin, M., Cardinal, A., et Bérard-Therriault, L. *Ardissonia crystallina* (Bacillariophyceae) : une étude en microscopie électronique à balayage des éléments structuraux des valves, 2686
Powell, G.R. see Remphrey, W.R., 268
Pryor, A., and Boelen, M.G. A double-stranded RNA mycovirus from the maize rust *Puccinia sorghi*, 2380

Purdy, B.G. see Macdonald, S.E., 1726
Quinn, J.A. see Bell, T.J., 2243
Rajaei, H. Changements cytochimiques et ultrastructuraux des parois cellulaires de la pellicule du raisin, *Vitis vinifera*, durant la croissance et la maturation de la baie, 1343
Rajchenberg, M. see David, A., 202
Raju, N.B., Perkins, D.D., and Newmeyer, D. Genetically determined nonselective abortion of asci in *Neurospora crassa*, 1539
Rao, K.S., and Catesson, A.-M. Changes in the membrane components of nondividing cambial cells, 246
Ratcliffe, M.J., and Turkington, R. Vegetation patterns and environment of some alpine plant communities on Lakeview Mountain, southern British Columbia, 2507
Rauser, W.E., and Ackerley, C.A. Localization of cadmium in granules within differentiating and mature root cells, 643
Reddy, P.S. see Bhagat, N.R., 1245
Redhead, S.A. The Xerulaceae (Basidiomycetes), a family with sarcodimitic tissues, 1551
Redhead, S.A., and Nagasawa, E. ***Resinomycena japonica*** and ***Resupinatus merulioides***, new species of Agaricales from Japan, 972
Reeleder, R.D. see Mercier, J., 1633
Regnier, E. see Peron, J.Y., 72
Reich, P.B., Schoettle, A.W., Stroo, H.F., Troiano, J., and Amundson, R.G. Effects of ozone and acid rain on white pine (*Pinus strobus*) seedlings grown in five soils. I. Net photosynthesis and growth, 977
Reid, C.P.P. see Cline, M.L., 869
Reid, D.M. see Macdonald, S.E., 1726
Reid, J., and Booth, C. *Winterella*, the correct name for *Cryptospora* and *Ophiovalsa*, 1320
Rembur, J. see Nougarède, A., 907
Remphrey, W.R., and Powell, G.R. Crown architecture of *Larix laricina* saplings: an analysis of higher order branching, 268
Remphrey, W.R., Davidson, C.G., and Blouw, M.J. A classification and analysis of crown form in green ash (*Fraxinus pennsylvanica*), 2188
Renaudin, S. see Benharrat, H., 1746
Rey, L. see Benharrat, H., 1746
Richardson, E.A. see Mims, C.W., 1236
Ringius, G.S., and Chmielewski, J.G. Morphological variation within and among six populations of *Trillium erectum* in southern Ontario, 2450
Ringius, G.S., and Semple, J.C. Cytogeography of the *Solidago spathulata – glutinosa* complex (Compositae: Astereae), 2458
Ringius, G.S. see Chmielewski, J.G., 1045
Ringius, G.S. see Chmielewski, J.G., 1671
Ritchie, J.C., Hadden, K.A., and Gajewski, K. Modern pollen spectra from lakes in arctic western Canada, 1605
Rittinger, P.A., Biggs, A.R., and Peirson, D.R. Histochemistry of lignin and suberin deposition in boundary layers formed after wounding in various plant species and organs, 1886
Roberson, R.W. see Mims, C.W., 1236
Robert, J.-C., and Bret, J.-P. Release of an inhibitor of stipe elongation from illuminated caps of *Coprinus congregatus* mushrooms, 505
Roelfs, A.P. see Al-Kherb, S.M., 994
Rondet, P. see Nougarède, A., 907
Rothwell, R.L. see Lieffers, V.J., 817
Routledge, R.D. Rhizome architecture for dispersal in *Eleocharis palustris*, 1218
Royer, J. see Foos, K.M., 1063
Rucker, C.J., and Zachariah, K. The influence of bacteria on trap induction in predacious hyphomycetes, 1160
Rybak, I. see Rybak, M., 2425
Rybak, M., and Rybak, I. Fossil chrysophycean cyst flora in a small meromictic lake in southern Ontario, and its paleoecological interpretation, 2425
Sack, F.D. The structure of the stem endodermis in etiolated pea seedlings, 1514
Sackston, W.E. see Gray, A.B., 2057
Saini, H.S., and Spencer, M.S. Manipulation of seed nitrate content modulates the dormancy breaking effect of ethylene on *Chenopodium album* seed, 876
Saini, H.S. see Goudey, J.S., 849
Sakai, S. Patterns of branching and extension growth of vigorous saplings of Japanese *Acer* species in relation to their regeneration strategies, 1578
Sampath, G. see Suryanarayanan, T.S., 815
Sanchez-de-Jiménez, E. see Loza-Tavera, H., 607
Sawhney, V.K. see Chandra Sekhar, K.N., 215
Sawhney, V.K. see Polowick, P.L., 807
Saxena, P.K., and King, J. Optimal conditions for the isolation and culture of protoplasts from a cell suspension culture of an isoleucine – valine-requiring auxotroph of *Datura innoxia*, 1736
Scagel, R. see Maze, J., 1531
Scagel, R.F. see Lindstrom, S.C., 2202
Scagel, R.K., El-Kassaby, Y.A., and Maze, J. Multivariate variation within and between open-pollinated families of Douglas-fir (*Pseudotsuga menziesii*), 1614
Schechter, S.E., and Gray, L.E. Oospore germination in *Phytophthora megasperma* f.sp. *glycinea*, 1465
Scheiner, S.M., and Teeri, J.A. Rapid genotypic change in a population of the grass *Danthonia spicata* following disturbance, 1819
Schoettle, A.W. see Reich, P.B., 977
Schoulties, C.L. see El-Gholl, N.E., 1733
Schubert, A. see Bonfante-Fasolo, P., 539
Schulte, P.J., Hinckley, T.M., and Stettler, R.F. Stomatal responses of *Populus* to leaf water potential, 255
Seastedt, T.R. see Hayes, D.C., 787

Seifert, K.A. ***Charomyces***, a new genus of Hyphomycetes with thallic–arthric conidia, 230
Seifert, K.A. *Stromatographium* and *Acrostroma* gen.nov.: two tropical hyphomycete genera with distinctive synnema anatomies, 2196
Seifert, K.A., and Carpenter, S.E. ***Bisporella resinicola*** comb.nov. and its ***Eustilbum*** anamorph, 1262
Sell, Y., et Cremers, G. Les inflorescences des Melastomataceae guyanaises, leur filiation et leur valeur taxonomique, 999
Semple, J.C., and Chmielewski, J.G. Revision of the *Aster lanceolatus* complex, including *A. simplex* and *A. hesperius* (Compositae: Astereae): a multivariate morphometric study, 1047
Semple, J.C. see Chmielewski, J.G., 1045
Semple, J.C. see Ringius, G.S., 2458
Serrano, B. see Loza-Tavera, H., 607
Shain, L., and Järlfors, U. Ultrastructure of eastern cottonwood clones susceptible or resistant to leaf rust, 1586
Sharik, T.L. see Holthuijzen, A.M.A., 1092
Sharik, T.L. see Johnson, W.C., 1622
Shay, J.M. see Zwiazek, J.J., 2310
Shimizu, S. see Kato, M., 729
Shipley, B. see van Hulst, R., 2373
Shoemaker, R.A., and Babcock, C.E. *Wettsteinina, 373*
Shoemaker, R.A., and Babcock, C.E. Additions to *Nodulosphaeria*, 1921
Shokes, F.M. see Ploetz, R.C., 2104
Siegel, M.R., Järlfors, U., Latch, G.C.M., and Johnson, M.C. Ultrastructure of *Acremonium coenophialum, Acremonium lolii*, and *Epichloe typhina* endophytes in host and nonhost *Festuca* and *Lolium* species of grasses, 2357
Sigal, L.L. see Eversman, S., 1806
Sigler, L. see Currah, R.S., 2473
Sihachakr, D., et Ducreux, G. Isolement et culture de protoplastes à partir de petioles et de tiges de deux variétés de patate douce (*Ipomoea batatas*), *192*
Simpson, S.J. see Owens, J.N., 1439
Sinclair, J., and Macdonald, P. Photosystem II activity and triazine resistance in weeds, 2147
Singh, A.P. see Harris, J.M., 280
Small, E. A taxonomic study of the "medicagoid" *Trigonella* (Leguminosae), 1199
Smart, M.G., Aist, J.R., and Israel, H.W. Some exploratory experiments on the permeability of papillae induced in barley coleoptiles by *Erysiphe graminis* f.sp. *hordei*, 745
Smereka, K.J., Machardy, W.E., and Kausch, A.P. Cellular differentiation in *Venturia inaequalis* ascospores during germination and penetration of apple leaves, 2549
Smith, D.H. see Mitchell, J.K., 2263
Smith, E.P. see Johnson, W.C., 1622
Smith, F.A. see Tester, M., 419
Smith, P.G., and Phipps, J.B. Studies in *Crataegus* (Rosaceae, Maloideae). XVI. Taxonomy of *C.* series *Rotundifoliae* in Ontario, 2646
Smith, S.E. see Tester, M., 419
Smith, W.H. see Coleman, J.S., 949
Sohlberg, E.H., and Bliss, L.C. Responses of *Ranunculus sabinei* and *Papaver radicatum* to removal of the moss layer in a high-arctic meadow, 1224
Solbreck, C., and Andersson, D. Vertical distribution of fireweed, *Epilobium angustifolium*, seeds in the air, 2177
Souvre, A. see Gélie, B., 2061
Spear, R.N. see Stanosz, G.R., 2124
Spence, J.R. see Ireland, R.R., 859
Spencer, M.S. see Goudey, J.S., 849
Spencer, M.S. see Saini, H.S., 876
Spicer, K.W. see Catling, P.M., 750
Sprent, J.I. see Faria, S.M. de, 553
Sreenivasa, B.A. see Wein, R.W., 1180
Standley, L.A. Taxonomy of the *Carex lenticularis* complex in eastern North America, 673
Stanosz, G.R., Patton, R.F., and Spear, R.N. Structure of *Armillaria* rhizomorphs from Wisconsin aspen stands, 2124
Stanton, M.L. see Nakamura, R.R., 2463
Staub, J.E., Fredrick, L., and Marty, T.L. Electrophoretic variation in cross-compatible wild diploid species of *Cucumis*, 792
Stefani, A., Meletti, P., and Onnis, A. Morphological characteristics of the experimental allopolyploid *Triticum durum* × *Haynaldia villosa* ($2n = 42$), 1948
Steijlen, I., and Zackrisson, O. Long-term regeneration dynamics and successional trends in a northern Swedish coniferous forest stand, 839
Stein, W.E., Jr. see Beck, C.B., 348
Stephen, F.M. see Paine, T.D., 2090
Stephen, F.M. see Paine, T.D., 2093
Stephens, L.C. see Hicks, C.B., 1967
Stettler, R.F. see Schulte, P.J., 255
Stevenson, S.C., and Lee, P.F. Ecological relationships of wild rice, *Zizania aquatica*. 6. The effects of increases in water depth on vegetative and reproductive production, 2128
St-Laurent, L., and Lalonde, M. Isolation and characterization of *Frankia* strains isolated from *Myrica gale*, 1356
Stockey, R.A. see Crane, P.R., 2490
Stone, J.K. Initiation and development of latent infections by *Rhabdocline parkeri* on Douglas-fir, 2614
Stroo, H.F. see Reich, P.B., 977
Strullu, D.G. see Bourgeais, P., 1989

Suryanarayanan, T.S., Muruganandam, V., and Sampath, G. Effect of Congo red on hyphal morphogenesis and sporulation of *Botryodiplodia theobromae*, 815
Suske, J., and Acker, G. Internal hyphae in young, symptomless needles of *Picea abies*: electron microscopic and cultural investigation, 2098
Szécsi, Á., and Tóth, S. Comparison of isolates in the *Chaetomium aureum* species group by esterase isoenzyme analysis, 198
Ta, T.C., Macdowall, F.D.H., and Faris, M.A. Utilization of carbon from shoot photosynthesis and nodule CO_2 fixation in the fixation and assimilation of nitrogen by alfalfa root nodules, 2537
Taber, R.A. see Mitchell, J.K., 2263
Tabor, C.A., and Barnett, N.M. An experimental system for studying interrelationships between the embryo and megagametophyte of *Pinus strobus* during seed germination, 1212
Talbot, S.S. Bryophytes from Nahanni National Park and vicinity, Northwest Territories, Canada. III, 592
Tam, C.E., Cole, K.M., and Garbary, D.J. *In situ* and *in vitro* studies on the endophytic red algae *Audouinella porphyrae* and *A. vaga* (Acrochaetiales), 532
Teeri, J.A. see Scheiner, S.M., 1819
Tester, M., Smith, S.E., and Smith, F.A. The phenomenon of "nonmycorrhizal" plants, 419
Thai, K.M. see Jana, S., 1741
Thain, J.F. see Côté, R., 509
Thalouarn, P. see Benharrat, H., 1746
Thériault, A. see van Hulst, R., 2373
Thompson, B.K. see Warwick, S.I., 1396
Thompson, B.K. see Warwick, S.I., 1403
Thorn, R.G. see Barron, G.L., 774
Tolonen, K. see Wein, R.W., 1180
Torrey, J.G. see Newcomb, W., 80
Tóth, S. see Szécsi, Á., 198
Trappe, J.M. see Hunt, G.A., 438
Trappe, J.M. see Carpenter, S.E., 716
Traquair, J.A. Backscattered electron imaging as a tool for histochemically localizing calcium oxalate with the scanning electron microscope, 888
Traquair, J.A. Mating incompatibility and cultural distinctions between *Coprinus phaeosporus* and *C. psychromorbidus*, 928
Traquair, J.A. Oxalic acid and calcium oxalate produced by *Leucostoma cincta* and *L. persoonii* in culture and in peach bark tissues, 1952
Traquair, J.A., Gaudet, D.A., and Kokko, E.G. Ultrastructure and influences of temperature on the *in vitro* production of *Coprinus psychromorbidus* sclerotia, 124
Tripathi, B.K. see Ammar, S., 137
Trites, L.F., and Bidwell, R.G.S. Effect of acidic precipitation on bean plants, 1121
Troiano, J. see Reich, P.B., 977
Turkington, R. see Aarssen, L.W., 1364
Turkington, R. see Ratcliffe, M.J., 2507
Turner, N.J., Kuhnlein, H.V., and Egger, K.N. The cottonwood mushroom (*Tricholoma populinum*): a food resource of the Interior Salish Indian peoples of British Columbia, 921
Tyrrell, L.E., and Boerner, R.E.J. *Larix laricina* and *Picea mariana*: relationships among leaf life-span, foliar nutrient patterns, nutrient conservation, and growth efficiency, 1570
Vallade, J., Maizonnier, D., et Cornu, A. La morphogenèse florale chez le pétunia. I. Analyse d'un mutant à corolle staminée, 761
Vallejos, C.E., and Pearcy, R.W. Differential acclimation potential to low temperatures in two species of *Lycopersicon*: photosynthesis and growth, 1303
Valmaseda, M., Martínez, A.T., and Barrasa, J.M. Annellidic conidiogenesis in *Pithoascus schumacheri* and redefinition of *Pithoascus* and related fungi, 1802
VandenBosch, K.A. see Hostak, M.S., 1108
Van den Heuvel, J. see Leone, G., 2133
van der Aa, H.A. see de Nooij, M.P., 2318
Vander Kloet, S.P., and Lyrene, P.M. Self-incompatibility in diploid, tetraploid, and hexaploid *Vaccinium corymbosum*, 660
van der Meer, J.P. Experimental hybridization of *Palmaria palmata* (Rhodophyta) from the northeast and northwest Atlantic Ocean, 1451
van Hulst, R., Shipley, B., and Thériault, A. Why is *Rhinanthus minor* (Scrophulariaceae) such a good invader?, 2373
Van Staden, J. see Mooney, P.A., 622
Van Tyne, C. see Wasserman, J.L., 2622
Vavasseur, A. see Lascève, G., 1861
Verma, R.K. see Kamal, 1259
Vieth J. see Dubuc-Lebreux, M.-A., 261
Vieth, J. see Laliberté, S., 299
Vieth, J. see Laliberté, S., 324
Vieth, J. see Dubuc-Lebreux, M.-A., 2074
Vincent, G. see Barabé, D., 1756
Visser, S., Parkinson, D., and Hassall, M. Fungi associated with *Onychiurus subtenuis* (Collembola) in an aspen woodland, 635
Vitt, D.H. see Glime, J.M., 1824
Voesenek, L.A.C.J., and Blom, C.W.P.M. Rooting patterns of *Rumex* species under drained conditions, 1638
Volkmann-Kohlmeyer, B. see Kohlmeyer, J., 571
Vose, J.M., and White, A.S. Processes of understory seedling recruitment 1 year after prescribed fire in an Arizona ponderosa pine community, 2280
Wagner, J. Ontogénie des périthèces d'*Endocarpon pusillum* (Pyrénolichen, Verrucariacées), 2441
Wagner, J. see Boissière, J.-C., 1468

Wainman, N., and Mathewes, R.W. Forest history of the last 12 000 years based on plant macrofossil analysis of sediment from Marion Lake, southwestern British Columbia, 2179
Walker, D.A. Height and growth rings of *Salix lanata* sp. *richardsonii* along the coastal temperature gradient of northern Alaska, 988
Wallen, D.G., and Allan, R. Utilization of amino acids by the blue-green alga *Synechococcus AN* (*Anacystis nidulans*), 1133
Wardenaar, E.C.P. A new hand tool for cutting peat profiles, 1772
Warne, T.R., and Lloyd, R.M. Gametophytic density and sex expression in *Ceratopteris*, 362
Warwick, S.I., and Thompson, B.K. Differential response to competition in weedy biotypes of proso millet, 1403
Warwick, S.I., Thompson, B.K., and Black, L.D. Life history and allozyme variation in populations of the weed species *Setaria faberi*, 1396
Wasserman, J.L., Mineo, L., Majumdar, S.K., and Van Tyne, C. Detection of heavy metals in oak mycorrhizae of northeastern Pennsylvania forests, using x-ray microanalysis, 2622
Weigle, J.L. see Hicks, C.B., 1967
Wein, R.W., Burzynski, M.P., Sreenivasa, B.A., and Tolonen, K. Bog profile evidence of fire and vegetation dynamics since 3000 years BP in the Acadian Forest, 1180
Weinberger, P., DeChacin, C., and Czuba, M. Effects of nonyl phenol, a pesticide surfactant, on some metabolic processes of *Chlamydomonas segnis*, 696
Wetherbee, R., Hill, D.R.A., and Brett, S.J. The structure of the periplast components and their association with the plasma membrane in a cryptomonad flagellate, 1019
Wetzel, R.G. see Carlton, R.G., 1031
Wheeler, N.C., and Guries, R.P. A quantitative measure of introgression between lodgepole and jack pines, 1876
Wheelwright, N.T. see Jones, E., 2291
White, A.S. see Vose, J.M., 2280
Whitecross, M.I. see Hawkins, C.D.B., 666
Whitney, H.S., Bandoni, R.J., and Oberwinkler, F. ***Entomocorticium dendroctoni*** gen. et sp. nov. (Basidiomycotina), a possible nutritional symbiote of the mountain pine beetle in lodgepole pine in British Columbia, 95
Whittier, D.P., and Given, D.R. The germination of *Tmesipteris* spores, 1770
Wicklow, D.T. see Frisvad, J.C., 765
Widholm, J.M. see Zehr, B.E., 491
Wierzbicka, M. Lead translocation and localization in *Allium cepa* roots, 1851
Wilcox, D.A., and Andrus, R.E. The role of *Sphagnum fimbriatum* in secondary succession in a road salt impacted bog, 2270
Wilkinson, H.T. Association of *Trechispora alnicola* with yellow ring disease of *Poa pratensis*, 150
Williams, M.E. see Zehr, B.E., 491
Williams, N.D. see Freeman, T.P., 2118
Williams, P.G. see Milligan, M.J., 598
Williams, P.H. see Leung, H., 112
Williams, P.H. see Bosland, P.W., 2067
Willison, J.H.M. see Zhang, M.I.N., 710
Willison, J.H.M. see Zhang, M.I.N., 1774
Wolf, S.J., and McNeill, J. Cytotaxonomic studies on *Polygonum* section *Polygonum* in eastern Canada and the adjacent United States, 647
Wolfe, C.B., Jr. Studies in the genus *Xanthoconium* (Boletaceae). I. New species and a new combination, 2142
Wolff, S.L., and Jefferies, R.L. Morphological and isozyme variation in *Salicornia europaea* (s.l.) (Chenopodiaceae) in northeastern North America, 1410
Wolff, S.L., and Jefferies, R.L. Taxonomic status of diploid *Salicornia europaea* (s.l.) (Chenopodiaceae) in northeastern North America, 1420
Woods, A.M., and Beckett, A. Wall structure and ornamentation of the urediniospores of *Uromyces viciae-fabae*, 2007
Woods, A.M. see Beckett, A., 1998
Wulff, R.D. Effects of irradiance, temperature, and water status on growth and photosynthetic capacity in *Hyptis suaveolens*, 2501
Yang, K.C. Metabolic activity of living sapwood ray cells in *Picea mariana* expressed by a new indicator, 1775
Yeung, E.C., and Law, S.K. The formation of hyaline caudicle in two vandoid orchids, 1459
Young, J.P.W. see Gould, K.S., 406
Zachariah, K. see Rucker, C.J., 1160
Zackrisson, O. see Steijlen, I., 839
Zehr, B.E., Williams, M.E., Duncan, D.R., and Widholm, J.M. Somaclonal variation in the progeny of plants regenerated from callus cultures of seven inbred lines of maize, 491
Zhang, M.I.N., and Willison, J.H.M An improved conductivity method for the measurement of frost hardiness, 710
Zhang, M.I.N., and Willison, J.H.M. Erratum: An improved conductivity method for the measurement of frost hardiness, 1774
Zwiazek, J.J., and Shay, J.M. Flouride- and drought-induced structural alterations of mesophyll and guard cells in cotyledons of jack pine (*Pinus banksiana*), 2310